

Review Article



Impact of Cultivation Condition and Media Content on *Chlorella vulgaris* Composition

Yunes Panahi¹, Ahmad Yari Khosroshahi^{2,3}, Amirhossein Sahebkar⁴, Hamid Reza Heidari^{2,5*}

¹Chemical Injuries Research Center, Systems Biology and Poisoning Institute, Baqiyatallah University of Medical Sciences, Tehran, Iran.

²Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

³Department of Medical Nanotechnology, Faculty of Advanced Medical Science, Tabriz University of Medical Sciences, Tabriz, Iran.

⁴Biotechnology Research Center, Pharmaceutical Technology Institute, Mashhad University of Medical Sciences, Mashhad, Iran.

⁵Department of Pharmaceutical Biotechnology, Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran.

Article info

Article History:

Received: 15 Mar. 2019

Revised: 17 Apr. 2019

Accepted: 4 May 2019

published: 1 June 2019

Keywords:

- Microalgae
- *Chlorella vulgaris*
- Nutraceuticals
- Cultivation condition
- Media composition

Abstract

Microalgae are a source material in food, pharmacy, and cosmetics industries for producing various products including high-protein nutritional supplements, synthetic pharmaceuticals, and natural colors. A promising algal source for such productions is *Chlorella vulgaris* which contains a considerable protein content. Similar to other microalgae, its desirability is minimal nutrient requirements since they are unicellular, photosynthetic, and fast-growing microorganisms. Another propitious option to be produced by *C. vulgaris* is biodiesel, since it is rich in oil too. Besides, algal well thriving in presence of increased amount of carbon dioxide makes them a practicable alternative biofuel resource without some problems of the traditional ones. At the same time, *C. vulgaris* is also a promising source for nutraceuticals such as amino acids, vitamins, and antioxidants. This review aims to discuss the conditions need to be observed for achieving a favorable growth efficiency of the *C. vulgaris*, as well as targeted productions such as biomass, antioxidant, and biofuel. Additionally, different approaches to induce any specific production are also considered comprehensively.

Introduction

Microalgae are located all around the world found in both aquatic (mostly) and all terrestrial ecosystems. They may live either individually or establish a symbiotic communication with other living organisms.¹ Microscopic algae as photosynthetic eukaryotic microorganisms, though contains similar organelles to land-based plants such as chloroplasts and nucleus, yield more efficiently biomass. The reason is their higher performance in utilizing sunlight, CO₂, water, etc leading to their extremely higher growth rates. Other algae advantage over plants is their ability to be cultured in various aquatic environments with less need for fertilizers and pesticides leading to less waste and pollution.^{2,3}

Microalgae are being used in multiple industries including food, pharmacy, cosmetics, and environment (producing biofuels and treatment of wastewater).⁴ For example, the whole biomass of some algal species are used as protein-repository for valuable chemical agents (pigments, enzymes, additive for cosmetics) and nutritional supplements; some *Spirulina*-containing preparations are

used in wound cicatrization for their therapeutical traits; *Scenedesmus*-containing preparations are suggested for treating some skin conditions (e.g. eczema); and finally some algal-originated ingredients have been addressed to suppress the HIV virus.⁴⁻⁸ Microalgae can remove the pollutants such as carcinogenic chemicals from wastewater, however, dark color and high-fat content can inhibit their growth. They can also be effectively used in the treatment of different animal farm effluents.^{9,10}

The extremely high oil content of some microalgae strains has made them great candidates for producing biodiesel. Producing biodiesel from microalgae in lieu of plants (routinely sweet corn) for production subsides the concerns about reducing human food, animal feed, and other products from plants.

Despite the expensive production process, *Chlorella vulgaris* still remains of the most promising algal strains for oil production in bioreactors.^{4,8,11} *C. vulgaris* is a very resilient species of unicellular green algae classified as chlorophyta. More than half of its biomass typically consists of protein (51%-58%), and the remained consists

*Corresponding Author: Hamid Reza Heidari, Email: heidarihr@tbzmed.ac.ir

of carbohydrate (12%-17%), lipid (14%-22%), and other different valuable nutraceuticals such as vitamins, antioxidants, and trace elements.⁴

Microalgae cultivation

Numerous arrangements have been attempted to obtain the ideal microalgal growth. To grow microalgae in the large volumes, two approaches are taken majorly: the outdoor ponds with sunlight as photo-supplier, and the outdoor or indoor photo-bioreactors (PBRs) with electric light-suppliers. Naturally, specific demands such as species control and cell growth optimization can be provided more conveniently in artificial bioreactors.^{12,13} The set-up materials, light source, circulation system through the reactor, CO₂ supplying system, the relative amounts of different nutrients in the medium, the pH of culture media, ambient and internal temperature of the growth medium, etc. are of basic features required to be measured when designing a PBR.^{12,13} Figure 1 shows different affecting parameters on the *C. vulgaris* growth output.

Process technology

Applied phycologists divide the microalgae cultivation systems into open and closed systems. In an open cultivation configuration, the culture medium is freely exposed to the ambient air under direct lighting of the sun; while, in a closed system the culture is under ceiling (indoor) isolated from the direct sunlight.¹⁴

Open system

In open systems, the ambient temperature optimally should be above 15°C and continuously not less than the freezing temperature. Also, the pond's depth should be adjusted between the light availability and the stirring optimality. The shallower bioreactors provide better light availability for algae cells, while adequate mixing requires a minimum depth. On the other hand, in shallow cultures, easy surface evaporation results in too much ionic fluctuation which is a negative factor for the algal growth.¹⁵ Therefore, the best dimensions of the bioreactor should be measured. Most

ponds are 20-30 cm in depth. Usually, the light availability becomes a limiting factor for the cell growth typically keeping the biomass concentration between 0.1 and 0.5 g/L.

The open systems are simple to design in the forms of lakes, tanks, and various pond types including those are open, covered open, shallow artificial (raceway) ones, shallow big ponds, circular ponds, etc.^{16,17} However, majorly the three designs of circular ponds, raceway ponds, and inclined ponds are upscaled for mass cultivation of microalgae.¹⁸ The disadvantage of open ponds is that only few extremophile or fast-growing algae (such as *Dunaliella* and *Spirulina*) can be grown productively in such systems. Because, only these species can conquer the harsh growth conditions existing in such systems (e.g. alkaline pH) and survive contamination.¹⁹ However, the contamination risk in open pond systems by other microalgal, fungal, bacterial, or protozoan species is irrefutable. In spite of its less cultivation efficiency, open ponds constitute the majority microalgae production around the world because of their feasibility in setting up.

Closed systems or photo-bioreactors (PBRs)

For the specific algal species with especial cultivation needs, fully closed systems structured from a transparent material called PBRs are better choices. PBR system provides controlled conditions such as temperature, culture density, pH levels, aeration rate, stirring program, carbon dioxide supply, water, and the light regime, around and in the culture medium. The main parameter about PBR is the best surface-to-volume ratio for highest light access in order to provide the most photosynthetic efficiency.²⁰

PBRs are commonly structured as flat plates or tubular reactors in which the water and nutrients are continuously flowed into the PBR.²⁰ The high surface-to-volume ratio in flat plate reactors provides the efficient use of sunlight.²¹ In these systems, narrow panels aligned in horizontal, vertical, or other angles in order to provide a uniform exposure to the light and air/CO₂ are supplied.²⁰ These




Figure 1. The main influential parameters on the efficiency and growth rate of *C. vulgaris* and its significant products are presented schematically in a single cell view.

systems, generally, purvey relatively condensed biomass and improved productivities.²⁰ Tubular reactors are often constituted of transparent tubes in the form of parallel loops arranged vertically, horizontally, or helically,²¹ and generally do not exceed one decimeter in diameter.²² The advantage of this system is that the culture medium is entirely separated from the environment to be preserved from the outer contaminants. CO₂ is pumped into the medium through a longitudinal laminar flow guaranteeing enough distribution and blending in order to attain up to 6 g/L of biomass. In return, the tubes cannot be too long because the nature of water does not allow the CO₂ or air to efficiently exchange between the environment and substrate for a long time. Then, high amounts of dissolved oxygen, sub-optimum CO₂ supplies, and increased pH levels are remained obstacles en route of the scale-up procedure of tubular PBRs.²¹

In a comparative study on different systems of *C. vulgaris* cultivation, Lam et al reported that this algae grow most efficiently in the presence of inorganic nutrients under indoor cultivation conditions.²³

Controlling of *Chlorella vulgaris* productions

Controlling physical parameters

A temperature adjustment according to the other algal species, usually suffices the *C. vulgaris* to survive. A cooling system should be considered for the temperature increase due to daylight to preserve an optimum growth rate.^{22,24} Sufficient stirring is required for good growth of algae and preventing the biomass sedimentation which is often provided by air/gas bubbling or pump-turbulating of culture medium. Mixing is especially important in reactors with a continuous light source to shift the cells from the dark to the illuminated volume for achieving a high productivity.¹ Ideally, the growth rate of microalgae is in a direct correlation with the rate of photosynthesis. Photosynthesis rate in the PBRs depends on the amount of solar radiation that technically cannot always penetrate all throughout the culture, so that the outer layers receive more light compared to the inner volume. In addition to light concentration and distribution, its kind (sunlight or artificial light) is also important to be considered in designing PBRs. Additionally, a dark period is required to be planned for the best result.¹

Light is defined as an electromagnetic radiation with wavelengths of between 10⁻³ and 10⁻⁸ meters ranging from radio waves and γ, to X-rays, and ultraviolet and visible lights. Among all, just the visible light (400-750 nm) can supply the required energy for the photosynthetic reactions.²⁵ At the same time, it can also be a limiting factor if the photons either dissipate as heat or reflects instead of being absorbed. The limitation in partial absorbing photon flux density originates from the limited optical properties of the cell or the culture mixing rate. While reflection happens to a small fraction of the photons, microalgal cell density increase exponentially until any

photons have remained absorbable for photosynthesis. Then, the cell growth remains constant until other limiting factors such as light density or nutrients level fall down.¹ Another important factor in designing bioreactors for microalgae is the light periods duration (photoperiods) that importantly affect on the microalgae's growth and photosynthesis efficiency. For example, the overexposure to light may cause an unnecessary overconsumption of electricity beside subsiding the cell growth. Whereas, a comparative study by Malaysian researchers showed a photoperiod of 12 hours (near local outdoor conditions) yielded much more microalgae biomass concentration compared to 3 h, 6 h, and 9 h photoperiods.²³ This proportional correlation between microalgae growth rates and light period duration remains the same at different light intensities.^{26,27} Higher productivities and maximum cell density values at 12:12h (light:dark) photoperiod, in comparison with the 14:10 h (light:dark)²⁶ shows that the overexposure to light reverses the results. On the other hand, a later study has shown that flashing lights could significantly increase the total fatty acid (TFA) content of *C. vulgaris* in addition to its growth rate either the lighting period was continuous or cycling (light-dark).²⁸ On the other hand, Zhao and colleagues showed that red light wavelength at the intensity of 1200-1600 μmol/m²/s is optimum for microalgal growth.²⁹ More studies regarding light impact on microalgae growth and production are summarized in Table 1.

The biomass concentration is also under the influence of initial culture pH parameter. High pH level can reduce the growth rate and the lipid output of some microalgal species, however, extremely higher pH can precipitate calcium salts.^{4,59} According to the reports, different strains of *C. vulgaris* tolerate low pH levels (from 3.0)^{60,61} better than high levels of pH (i.e. 11).⁵⁹ In this regard, pH=2 stops *C. vulgaris* growth completely, pH=4 leads in reduced cell number, while pH=6 cause that *C. vulgaris* grow faster than in pH=8 (however, they yield similar final cell number), and higher pH (10 and 12) reduce growth even more. Then, apparently, pH=6 is the approximate desired pH for *Chlorella* cultivation.⁶²

Controlling media components

The growth level as well as the final concentration of microalgae are highly influenced by media composition.⁶¹ Nutrient-rich (eutrophic) waters lead to more microalgal blooms.⁶³ Providing all essential nutrients in the growth medium of algae is necessary for adequate biomass production.²⁴ The first and the most important ingredient that contributes to the algal production is carbon. Organic sources of carbon such as acetic acid or peptone as well as inorganic from of carbon dioxide can be supplies in the medium.¹ Nitrogen is the next essential nutrient involved in algal biomass production usually added to the medium in the nitrate (NO₃⁻) form.^{1,64} Nitrogen consists more than 10% of the biomass content and its shortage in culture

Table 1. The effects of different conditions and compositions on microalgal growth for achieving the highest amount of biomass, protein, antioxidants, lipids, and carbohydrates

Parameter	Mechanism/Strategy	Achievement	Ref
Protein and Antioxidant			
NO ₃	More nitrate utilization and more protein accumulation	Protein content increased up to 44.3%	30
Traumatic acid (10 ⁻⁶ -10 ⁻⁵ M)	Increase in antioxidant enzymes activity (SD, catalase, POD, GR)	The lipid peroxidation stopped by TA SH-group proteins underwent oxidative destruction	31
N-phenyl-2-naphthylamine (2.5 mg/L)	Inhibited photosynthesis, triggered ROS synthesis, disrupted the subcellular structure	As an allelochemical, 2.5 mg/L of it significantly increased antioxidant enzymes activities e.g. SOD, POD, and catalase	20,32
Nonylphenol (0.1-1.0 mg/L) exposure time	Induced oxidative stress	Obvious effects on antioxidant responses in the first day High NP contents changed the SOD, catalase, POD, and GSH levels	20,27,33
Elevated light intensities (400 µmol photon/ms)	Potential source of zeaxanthin ^a So-called "molecular sunglasses" mechanism	Induced color change of microalgae from green to yellow Extra produced xanthophylls act as potential antioxidants, stabilize the membrane, and protect cells from intensive radiation	34
Iron-dependent oxidative stress	Triggered oxidative stress by surplus iron, decreasing the cellular growth rate of phytoplankton	>200 µM iron supply reduced the <i>C. vulgaris</i> growth level and did not changed the β-Carotene content >90 µM iron availability raised vitamin E ^b , vitamin C ^b , and total thiol content	35
Sodium Nitroprussiate CTAC/Flu surfactant Polycyclic aromatic hydrocarbons	SNP alleviated the pollution damage of surfactants and PAHs by providing external NO ⁻ for <i>C. vulgaris</i> cells	Supplying 20 µM SNP: Increased the biomass, the chlorophyll concentration, and the activity of SPC, SOD, POD, and catalase Decreasing MDA and ROS amounts	20,36
Trifloxystrobin	Decreased antioxidant enzymes' activity Disturb photosynthesis in <i>C. vulgaris</i> Destruct the cellular structure	255.58 µg/L of trifloxystrobin (IC ₅₀): Reduced transcription of genes associated with the photosynthesis, soluble proteins, and T-AOC Increased SOD & POD activities and ATP expression	37
Azoxystrobin	AZ disrupts the <i>C. vulgaris</i> growth through: reducing energy/photosynthesis-associated mRNA expressions Inducing ROS overproduction	510 µg/L (IC ₅₀) of AZ: Reduced the chlorophyll and soluble protein content Increased the T-AOC level Weakened SOD, POD, GSTs, and GPx activities, and GSH content	38,39
Lipid and Biomass			
Novel lighting methods	Persistent illumination, periodical light-dark durations, persistent darkness with additional flashing light, complete darkness	The flashing light increased the growth rates of <i>C. vulgaris</i> significantly and TFA concentrations	28
Nitrogen and some trace elements availability	Utilized N, Mn, Ca, etc.	Lipid production reached approximately threefold	40
N and P content optimization	N concentration ranged 0–56 mg/L and P ranged 0–19 mg/L	N/P ratio =10: At the end of cultivation, P and N were totally eliminated and biomass concentration had reached 1.58 g/L	41
Light limitation Nitrogen starvation	By external shading and dimming the algal bags	Algal growth reduced due to light-limitation and nitrogen starving For future studies, separate analysis of biomass and lipid yield is suggested	42

Table 1. Continued

Parameter	Mechanism/Strategy	Achievement	Ref
Nitrogen source limitation (ammonium/nitrate)	Optimization study of sodium nitrate, ammonium bicarbonate, heptahydrate magnesium sulfate, potassium dihydrogen phosphate, dipotassium phosphate, diammonium phosphate Based on BBM and HAMGM	Optimizing the culture medium improved: Biomass: 40% (0.73 g/L) regarding BBM Lipid concentration: 85% (281 mg/L) regarding HAMGM	43
Harvest time optimization of chlorella growth factor	Harvest time effect of CGF extract as a growth stimulator	CGF enhanced the lipid and total biomass levels (>1 kg/m ³) after 5 days	44
Temperature Nitrogen concentration Initial cell density	Temperature ranged 20-30°C	The optimal temperature for the neutral lipid productivity: 27-27.4 °C Optimal nitrogen concentration for the algal growth: 1.5 g/L Optimal cell density: 50%	45
CO ₂ concentration Light intensity Nitrogen deficiency Phosphorus luxury uptake	CO ₂ : 0.03, 4, 6, 12 percent Light intensity: 40, 120, 200 µmol photons/m/s) Nitrogen deficiency combined with pre-optimized phosphorus	4% CO ₂ concentration and light intensity of 200 µmol photons/m/s provided the best results Polyphosphate maximum uptake rate: 2.08 mg/L/day Better performance under nitrogen deficiency than nitrogen sufficiency	46
MgSO ₄ ·7H ₂ O, KNO ₃ , Glucose, and NaCl	Using the Plackett–Burman design to selected the key nutrient factors Applying the Box–Behnken design to optimize the strategy	Maximum predicted biomass concentration (4.28 g/L) was verified with these calculated amounts: glucose (25 g/L), MgSO ₄ ·7H ₂ O (1.33 g/L), KNO ₃ (1.30 g/L), NaCl (3.02 g/L)	47
Glucose, acetate, and glycerol	Autotrophic, heterotrophic, and mixotrophic modes	The heterotrophic nutrition mode yielded the maximum biomass (8.9 g/L) and lipid contents (36.19%) in CM	48
Red light intensity	Optimization Red light intensities: 800, 1200, 1600, and 2000 µmol/m ² /s	The best visible light for the optimized microalgae growth was red wavelength The optimal concentration of light was 1200-1600 µmol/m ² /s	29
Inorganic carbon, phosphorous Light intensity	Chemostat cultivation	Maximum biomass (538 mg/L/d) and lipid (128 mg/L/d) contents achieved at the first phase	49
Phytohormones	Auxins and gibberellins range 40-60 µM Microalgae spp: <i>Scenedemus abundans</i> , <i>Chlorella ellipsoidea</i>	Phytohormones promoted the microalgae due to reducing the intracellular ROS level >3-fold increment for <i>S. abundans</i> ~7-fold increment for <i>C. ellipsoidea</i>	50
Blue green-11 medium, Bold basal medium, Fog's medium, and Basal medium	Comparison screening Optimization	BG-11 yielded higher lipid content A2:1 mixture of chloroform:methanol was identified as the most effective ratio, extracting an average of 15% total lipids	51
Nitrogen sources: Urea, KNO ₃ , NaNO ₃ , and NH ₄ NO ₃	Optimization Microalgae Sp.: <i>Chlorella sorokiniana</i> Urea concentrations: 0-10 g/L	1.5g/L urea purveyed the highest biomass concentration (0.220 g/L) and 61.52% lipid content	52
Light condition light/dark photoperiod	Optimization	Suitable light intensity: 2000Lux Appropriate light-dark photoperiod: 12:12	53
Sodium Erythorbate as common antioxidant	NaE range: 2.0-16.0 g/L Moderate controlling the accretion of the dissolved photosynthetic oxygen in the media	Algal autotrophic growth enhanced effectively The NaE treatment provided higher pigment contents (4.17 to 4.44 times), cell density (2.67 times), and algal biomass (1.21 times) compared to the glucose treatment	54

Table 1. Continued

Parameter	Mechanism/Strategy	Achievement	Ref
Lipids and Carbohydrates			
Light concentration CO ₂ level Stirring	Hybrid bioreactors benefiting the features of both pneumatic bioreactors and stirred tanks	Microalgae grew exponentially for around one week Entered the stationary phase after 9-14 days	55
Carbon sources: Glucose Glycerol Acetate	Glucose optimization using mixotrophic cultivation Assessing the usability of wastewater for renewable biomass and high-value microalgal oil production	Appropriate carbon source concentration: 5 g/L glucose Optimized biomass yield: 0.13 g/L/d Dry cell weight: 1.39 g/L Total lipid content: 19.29 ± 1.83% Total carbohydrate: 41.4 ± 1.46%	56
Glucose Nitrate Phosphate	Artificial neural networks A feed-forward method Optimization	The optimized medium condition: glucose (15 g/L), N (1.04 g/L), P (0.005 g/L) The highest lipid yield: 1.944 g/L The highest biomass yield: 0.31 g/L/d	57
Light	Supplying glucose substrate for the oil-rich microalga <i>Chlorella zofingiensis</i>	Light attenuates lipid accumulation possibly by inhibiting lipid biosynthetic pathway and promotes proliferating the cells and starch synthesis Attenuated lipid accumulation caused carbon transform from glucose to starch	58

Abbreviations; SD: sodium dismutase, TA: Traumatic acid, POD: ascorbate/NADH peroxidase, GR: glutathione reductase, ROS: reactive oxygen species, SOD: superoxide dismutase, NP: Nonylphenol, GSH: glutathione, SNP: Sodium Nitroprussiate, PAHs: polycyclic aromatic hydrocarbons, SPC: soluble protein content, MDA: malondialdehyde, T-AOC: total antioxidant contents, AZ: Azoxystrobin, GSTs: glutathione S-transferase, GPx: glutathione peroxidase, TFA: total fatty acid, BBM: Bold's Basal Medium, HAMGM: Highly Assimilable Minimal Growth Medium, CGF: chlorella growth factor, CM: culture medium, BG-11: Blue green-11, medium, NaE: Sodium Erythorbate.

^a A macular pigment that protect eyes against age-associated macular degeneration

^b Vitamin E: α -tocopherol, vitamin C: ascorbate.

medium leads to decreasing the chlorophylls, increasing carotenoids, accumulation of polysaccharides and certain oils,¹ and increasing lipid production.⁶⁴ Nitrogen deprivation gradually reform the lipid constituents from free fatty acids-rich compositions to triglyceride-rich lipids.⁶⁴ The third essential element in the medium for the cellular processes involved in the growth such as DNA anabolism and energy transmission, is phosphorus. It is mostly added in the form of orthophosphate (PO_4^{3-}).¹ Other elements involving selenium, potassium, sodium, iron, magnesium, and calcium plus the trace elements such as boron, copper, manganese, zinc, and molybdenum are other important nutrients mostly used in enzyme reactions.¹ In general, the water used for preparing the cultivation systems are recommended to be distilled, filtered, or de-ionized. However, the groundwater or any other available domestic aqua is acceptable for making large-scale growth medium.¹

Although the best conformation of culture medium for growing *C. vulgaris* has remained controversial, blue green-11 (BG-11) has been the most common medium of choice.⁶⁵⁻⁶⁷ This culture medium mostly consists of NaNO_3 and K_2HPO_4 ⁶⁸ as supplying resources of nitrogen and phosphorus. However, other culture media providing enough nitrogen and phosphorus are also frequent.^{64,69}

There are other factors which should be specifically controlled during algae cultivation. The biggest is the saline concentration of the culture media. Excessive amounts of salt can inflict the growth and metabolism of algae cells and reduce the growth of *C. vulgaris*.^{70,71} Less than four decades ago, the positive effects of little amounts of NaHCO_3 and bubbling with CO_2 enriched air on the *C. vulgaris* have been approved.⁷² For example, a concentration of 3.36 g/L of NaHCO_3 could yield 0.67 g/L of dried biomass.⁷³ Recently, it has also been shown that ammonium in the form of NH_4Cl is firstly consumed by *C. vulgaris* before it goes for the nitrate present in the inoculum.⁶ For different strains of *Chlorella*, a trace amount of nitrates has been enough for obtaining adequate biomass and lipid concentration, so that even 1.24 g/L KNO_3 sufficiently nurture the cultivation.⁷⁴ A comparison of cultivating *Chlorella sorokiniana* in presence of either 15 g/L of ammonium chloride or 1.5 g/L of sodium nitrate, resulted in same biomass concentration (both in Tris-Acetate-Phosphate media bubbled with 5% CO_2 enriched air). However, in the NH_4Cl -containing culture medium, the pH decreased more radically.⁷⁵ Adding sodium bicarbonate to the basal media in a range of 0.1-1.6 g/L, showed that maximum biomass production (0.6 g/L) of *C. vulgaris* takes place at the 1.2 g/L concentration of sodium bicarbonate.⁵⁹ Also, fixation of dissolved CO_2 in the bicarbonate form using NaOH instead of adding CO_2 alone, has led to more increases the final harvested biomass of *Chlorella* spp.⁷⁶

Beside the total biomass, the production level of other microalgae's constituents such as proteins, lipid,

carbohydrates, and antioxidants is also under the influence of media content. For instance, Xie and colleagues showed that after a deprivation period, *C. vulgaris* consumed more nitrate and produced more proteins up to 44.3%.³⁰ In an optimization study, a combination of nitrogen, magnesium, calcium, and some other trace elements provided a synergistic multi-parameter boosting effect and tripled the lipid production.⁴⁰ Adding NaHCO_3 to the media along with bubbling with CO_2 enriched air also increases the lipid production of some microalgal species. Maximum concentration of lipid for *Chlorella* have been shown to happen at the 75 mg/L concentration of NaHCO_3 with 4758 ppm of CO_2 . In addition, a glucose concentration of 5 g/L in *C. vulgaris* culture medium significantly increased the ultimate harvested biomass containing higher amounts of lipid, carbohydrate, and proteins.⁷⁷

Furthermore, microalgae are addressed as source of nutraceutical antioxidants due to their rich content of carotenoids, vitamins, and phenolics. Various studies have aimed maximizing the yield of antioxidants by induction of different stress factors. As an example, Sun et al showed that trace amounts of selenium ($<75 \text{ mg L}^{-1}$) can increase *C. vulgaris* cell growth rate, organic selenium content, and antioxidant activity through increasing production of some photosynthesis associated pigments and heme containing enzymes such as guaiacol peroxidase (GPx), catalase (CAT), and superoxide dismutase (SOD). However, overexposure to the selenium ($>100 \text{ mg/L}$) can reverse these changes.⁷⁸ More elaboration on the antioxidant production by microalgae and important affecting factors will be discussed in following section. Besides, details of the best condition for microalgae total lipid content in regards to biofuel production will be reviewed later.

The highest amounts of *C. vulgaris* biomass, and its protein, antioxidant, lipid, and carbohydrates contents achieved through variation in cultivation conditions and media formulations are summarized in Table 1. The studies are categorized based on the main affected production of *C. vulgaris* and some other species of *Chlorella*.

Induction of antioxidant production in *C. vulgaris*

In the general metabolism process of green microalgae, like *C. vulgaris*, oxidant reactive molecules such as H_2O_2 , free radicals derived from molecular oxygen or chemically active oxygen species (ROS), and oxidized lipid derivatives produced at chloroplast, mitochondria and peroxisomes compartments which is reviewed before.⁷⁹ Meanwhile, these oxidizing agents were scavenged and deactivated through adapted Enzymatic and Non-Enzymatic antioxidant pathways (Figure 2).

Normally, there is a balance between production of oxidant and antioxidant molecules inside the cell; as if there was an augment in oxidizing agent production due to some stressful conditions, cell prevents the hazardous




Figure 2. The most important stressful factors which induce oxidant and following antioxidant level of microalgae presented schematically in a single cell view.

effect of accumulated oxidizing agents by overproducing antioxidant molecule.⁸⁰ Accordingly, numerous strategies have been adopted, in which by providing the artificial stressful cultivation condition, microalgae antioxidant machinery have been boosted up.⁸¹ We will discuss light intensity, temperature, osmotic stress and nutrient limitation as main approaches to enhance antioxidant production in the following sections.

Light

Light intensity considered as one of the most important antioxidant production inducer. While electron transfer apparatus in photosystems has been saturated, or NADP supply of the cell has been limited, photo-inhibition occurred and the excitation energy transferred to oxygen and produce reactive oxygen derivatives.⁸² In this stressful condition several adaptation mechanisms have been taken, which mainly include: 1) reducing the antenna size, 2) dissipating the energy as heat, 3) CO₂ fixation in the forms of storage materials such as starch and lipids to redeem the reductive pressure of the electron transport chain, and 4) quenching ROS by antioxidant production.⁸¹ The photo-inhibition and oxidants overproduction enhance while microalgae are under the stressful growth limiting condition due to nutrient limitation or other unfavorable cultivation condition. The light stress can also promote algae to produce antioxidants including carotenoids, some vitamins, and butylated hydroxytoluene (BHT).

As an example, 4.4 kJ/m dose of UV-B exposure was taken as a stressful condition in a study to increase antioxidant value of *C. vulgaris*. This approach significantly increased radical content in the membrane as well as the superoxide dismutase activity by 40% and that of catalase by 500%.⁸³ Moreover, Grudzinski and colleagues showed that intense lightening (400 μmol photons/m/s) induced yellowing of green microalgae, over-synthesizing xanthophylls. Xanthophyll is a carotenoid with potential to act as antioxidant, stabilize the membrane and, specially, protect cells from severe radiation, via “molecular sunglasses” mechanism.³⁴

Temperature

Low temperature is one of the methods to increase antioxidant level in microalgae. Algae produce unsaturated fatty acids adding them to the membranes to sustain its fluidity in order to conform with the cold environment. Other adaptation mechanisms contain a) producing cold shock and antifreeze proteins, b) producing more enzymes to counterbalance the poor specific activity of enzyme,⁸⁴ and c) involving differential energy partitioning.

Moreover, in higher temperature there are some evidences of increased production of astaxanthin,⁸⁵ unsaturated fatty acids, ice structuring proteins, and phenols in some microalgae species. Miranda et al showed almost doubling the antioxidant activity of *C. vulgaris* culture at 30°C. This is possibly due to the increase of different phenolic compounds in the methanolic extract.⁸⁶

Osmotic stress

Variation in salinity is one of the important reasons of osmotic changes which can impose hypo- or hyperosmotic tension on the microalgae cells. Osmotic adaptation involves (a) water flux, (b) ion transport, (c) producing vacuoles to isolate the redundant ions,⁸⁷ and (d) producing one or several organic osmolytes such as glycerol.^{88,89}

Salinity tolerance depends on environmental conditions such as light and temperature as well as nutrient constraint. Hyper-salinity sometimes increase the production of the metabolites (e.g. palmitic and oleic acids), as well as carotenoids (e.g. lutein, β-carotene),⁹⁰ or astaxanthin.⁸⁸ In the result of osmotic stress, valuable metabolites including glycerol, unsaturated fatty acids, and carotenoids may also increase in the microalgae cultivation. Pandit and colleagues in a recent study evaluated salinity stress (0.06 to 0.4 M NaCl) on *C. vulgaris* biomass and its productions. Their results confirmed salt tension as a significant stress that not only cause a fatty acid production enhancement and improve their composition, but also augment the antioxidant production level of *C. vulgaris*.⁹¹

Nutrient limitation

Deprivation of any major nutrients leads to the cessation

of microalgae growth cycle and changing other cell metabolites output. Thus, the microalgae takes different adaptation mechanisms including a) up-regulation of enzymes associated with the mechanisms of limiting nutrient absorption; b) decrease in photosynthetic activity; c) oxidative stress following to Calvin cycle inhibition; d) producing large amounts of carotenoids;⁹² e) significant increase of protective pigments such as astaxanthin mainly via key nutrient deficiency (e.g. N, P, or S deprivation)⁹³; f) increase cellular content of lipids (mostly taken as a response to nitrogen starvation),⁹⁴ and g) upgrading the amount of polyunsaturated fatty acid.⁹⁵

Therefore, heightening the producing level of secondary carotenoids, polyunsaturated fatty acid, and vitamin E⁹⁶ as important antioxidants can be induced by nutrient limitation in microalgae. In the same regard, specifically in *C. vulgaris*, lipid production efficiency is mostly induced by the sulfur deprivation. Sakarika et al achieved a high level of total lipid ($53.43 \pm 3.93\%$ gram per gram dry weight) in a kinetic study.⁹⁷

Induction of Biofuel production in *C. vulgaris*

Exponentially reduction of non-renewable energy resources has propounded microalgae as an alternative source for producing biodiesel. The sustainable photosynthetic efficiency of microalgae plus the eco-friendly approach of algal petrodiesel generation have promoted their potential. Also, the non-arable land is utilized for biodiesel production from algae which appease the food versus fuel dilemma. Biodiesel is considered as a non-toxic and sustainable fuel because of its biologic origin and degradability. Additionally, the gaseous pollutants emitted by biodiesel is confirmed to be less than fossil diesel and it does not show negative effect on the carbon dioxide or sulfur amounts of the atmosphere.⁶⁴ Furthermore, microalgae cultivation for biofuel production can be conjoined with wastewater which is reasonable and helpful in terms of economy and environment conservation. Using algae for biofuel production is a sustainable approach, since during which lots of water and nutrient (e.g. N and P elements) are recovered.⁴

Before, there have been some controversy on the successfulness of biofuels as sustainable alternative for energy resources. Because, biofuels were firstly generated from agricultural edible products (including sugarcane, sugar beet, and corn) that primarily overlapped with the staple food of people. Then, one would discuss that world's food and water markets could be at risk and it also could compromise the forests. Next, biofuels were suggested to be produced from lignocellulosic substrates reminded from agriculture and wood industries whose converting technology to liquid biodiesel is not commercially productizing enough. The third generation biofuels seem to be technically sustainable resolving many of previously addressed problems. This group of biofuels use algae as

their resources for energy production.⁹⁸ Compared to the soybeans and sunflowers original lipid content (20% and 55% respectively), the primary lipid content of algae is not considerably high (2%-40%); while its per-area-unit output of biofuel is substantially higher (>50-fold). Also, the extraction process is much more economic. The biofuel production efficiency of algae derives from its proficiency of photosynthesis, high growth rate, and the simple structure of the algae that makes its biomass much more utilizable in the biofuel making industry.^{64,67,69,75}

In contrast to the plants, microalgae store their primary energy in lipid materials instead of saccharides. Microalgal primary energy storage may be in any form of fatty acids or their derivatives or any other substances and compounds that is biosynthetically or functionally related to them. Then, lipids and fatty acids are found in not only the microalgal membrane, but also as the metabolites and mere storage bodies in the cytoplasm, often located around the mitochondria. The oil molecules in microalgae are majorly found as tri-alkyl glycerides (triple long-chain fatty acids bonded to an axis of glycerol). However, isoprenoids, phospholipids, glycolipids, and hydrocarbons are also found. The microalgal lipids are mostly methyl esters which are more oxidized and more viscous compared to the crude petroleum.^{64,67,69,75}

Chlorella vulgaris has a great potential for producing biofuel. Previously, Yoo et al showed that the maximum lipid concentration in *C. vulgaris* can reach about 11% of its final dry weight if aerated using 10% CO₂-enriched air, putting it far after *B. braunii* that could produce oils up to about 25% of its total dry biomass.⁶⁷ But, Widjaja et al displayed that lipid production of the *C. vulgaris* grows in the presence of further CO₂ (~30% of its dry biomass weight). As mentioned before, it has been shown that *C. vulgaris* lipid production can also exceed 50 percent of the algae dry weight through growing in high CO₂ levels under nitrogen-deprived condition.⁶⁴

Based on a 2010 report, ~1% of the total cultivable land on the earth (equivalent to 14 million hectares) is under plantation for supplying resources of the producing biofuels equivalent to 1% of the global -only- "transport" fuels.⁹⁸ While, if a given algae species produce oil as much as the 30 percent of its biomass, it can produce 1,535 kg biomass/m²/d, which provides enough biodiesel for the U.S oil-derived transport in a year, only on the 3% of U.S arable terrains.²²

The heating value of algae-derives biofuel is between 40-45 MJ/kg, similar to that of fossil petroleum. Most attempts for producing biodiesel from algae after extracting the algal oil are aimed at using the same trans-esterification process as any other oilseed for producing biodiesel. Trans-esterification happens during the conversion of triglycerides in the vegetable/animal oils or free fatty acids into fatty acid alkyl esters and glycerol. This response takes place in presence of alcohol, heat, and a strong alkaline catalyst. The procedure is a well-established biodiesel

production from both vegetable and algal oils.^{64,67,69,75}

Anaerobic fermentation of microalgae biomass is the most simple method for producing energy from the algae due to its independency from the biomass composition.⁹⁹ During anaerobic fermentation, the organic carbons in the biodegradable substrate are digested to the form of methane (so-called biogas) during a four-stages metabolism including 1) hydrolysis, 2) acidogenesis (producing volatile fatty acids), 3) acetogenesis (producing acetic acid from volatile fatty acids), and 4) methanogenesis (converting acetates into methane). Microalgae is among the different microorganisms contributing to biogas production with a high production efficiency.¹⁰⁰ The reason is the high lipid content of the microalgae and lacking lignin compared to the conventional substrate used for biofuel production.⁶³ The cell composition of microalgae significantly impacts on the yield of biofuel, since the lipids convert to oil more conveniently than proteins, and carbohydrates. Therefore, the high output of lipid and protein and the digestibility of the microalgal cell wall because of the absence of lignin bring microalgae feedstock as a highly suitable substrate producing methanolic biogas.⁶³

Conclusion

Considering the potentials of managing microalgal cell culture conformation and its fast growth, different photosynthetic algal species are addressed to be suitable resource for producing CO₂-free nutraceuticals and biofuels. However, at present, the commercial cultivation of microalgal biomass mainly relates to a small high-value sector of industries such as food supplements or cosmetic ingredients. Up-scaling of this process needs more efforts to make the commercial microalgae production of nutraceuticals and biofuels and the required facilities should be financially justifiable. Then, more investigations are required to further improve the stability and productivity of outdoor microalgae culture systems. The systems reviewed above, help optimization of key parameters effecting on biomass, protein, antioxidant and lipid (biofuel) production efficiency. Accordingly, parameters such as light availability, temperature and nutrient composition should comprehensively regarded based on the main product of industrial facilities. However, optimization on out-door culture systems is still a remarkable limitation to address.

Ethical Issues

Not applicable.

Conflict of Interest

There is no conflict of interest to declare.

References

- Cysewski GR, Lorenz RT. Industrial production of microalgal cell-mass and secondary products-species of high potential: *Haematococcus*. In: Richmond A, ed. *Handbook of Microalgal Culture: Biotechnology and Applied Phycology*. Blackwell Publishing Ltd; 2004:281-8. doi: 10.1002/9780470995280.ch14
- Gouveia L. *Microalgae as a feedstock for biofuels*. Springer; 2011:1-69. doi: 10.1007/978-3-642-17997-6
- Iwamoto H. Industrial production of microalgal cell-mass and secondary products-major industrial species: *Chlorella*. In: Richmond A, ed. *Handbook of Microalgal Culture: Biotechnology and Applied Phycology*. Blackwell Publishing Ltd; 2004:253-63. doi: 10.1002/9780470995280.ch11
- Becker EW. *Microalgae: Biotechnology and Microbiology*. Cambridge University Press; 1994.
- Aslan S, Kapdan IK. Batch kinetics of nitrogen and phosphorus removal from synthetic wastewater by algae. *Ecol Eng* 2006;28(1):64-70. doi: 10.1016/j.ecoleng.2006.04.003
- Feng Y, Li C, Zhang D. Lipid production of *Chlorella vulgaris* cultured in artificial wastewater medium. *Bioresour Technol* 2011;102(1):101-5. doi: 10.1016/j.biortech.2010.06.016
- Gellenbeck KW. Utilization of algal materials for nutraceutical and cosmeceutical applications--what do manufacturers need to know? *J Appl Phycol* 2012;24(3):309-13. doi: 10.1007/s10811-011-9722-z
- Gustafson KR, Cardellina JH 2nd, Fuller RW, Weislow OS, Kiser RF, Snader KM, et al. AIDS-antiviral sulfolipids from cyanobacteria (blue-green algae). *J Natl Cancer Inst* 1989;81(16):1254-8. doi: 10.1093/jnci/81.16.1254
- Zeng X, Guo X, Su G, Danquah MK, Zhang S, Lu Y, et al. Bioprocess considerations for microalgal-based wastewater treatment and biomass production. *Renew Sust Energ Rev* 2015;42:1385-92. doi: 10.1016/j.rser.2014.11.033
- Goncalves AL, Pires JCM, Simoes M. A review on the use of microalgal consortia for wastewater treatment. *Algal Res* 2017;24:403-15. doi: 10.1016/j.algal.2016.11.008
- Heredia-Arroyo T, Wei W, Ruan R, Hu B. Mixotrophic cultivation of *Chlorella vulgaris* and its potential application for the oil accumulation from non-sugar materials. *Biomass Bioenergy* 2011;35(5):2245-53. doi: 10.1016/j.biombioe.2011.02.036
- Xu L, Weathers PJ, Xiong XR, Liu CZ. Microalgal bioreactors: challenges and opportunities. *Eng Life Sci* 2009;9(3):178-89. doi: 10.1002/elsc.200800111
- Juneja A, Ceballos RM, Murthy GS. Effects of environmental factors and nutrient availability on the biochemical composition of algae for biofuels production: a review. *Energies* 2013;6(9):4607-38. doi: 10.3390/en6094607
- Tredici MR. Mass production of microalgae: photobioreactors. In: Richmond A, ed. *Handbook of Microalgal Culture: Biotechnology and Applied Phycology*. Blackwell Publishing Ltd; 2004:178-214. doi: 10.1002/9780470995280.ch9
- Borowitzka MA. Commercial production of microalgae: ponds, tanks, tubes and fermenters. *J Biotechnol* 1999;70(1-3):313-21. doi: 10.1016/S0168-1656(99)00083-8
- Mata TM, Martins AA, Caetano NS. Microalgae for biodiesel production and other applications: A review. *Renew Sust Energ Rev* 2010;14(1):217-32. doi: 10.1016/j.rser.2009.07.020
- Singh J, Gu S. Commercialization potential of microalgae for biofuels production. *Renew Sust Energ Rev* 2010;14(9):2596-610. doi: 10.1016/j.rser.2010.06.014
- Richmond A. Biological principles of mass cultivation.

- In: Richmond A, ed. *Handbook of Microalgal Culture: Biotechnology and Applied phycology*. Blackwell Publishing Ltd; 2004:125-77. doi: 10.1002/9780470995280.ch8
19. Pulz O, Scheibenbogen K. *Photobioreactors: design and performance with respect to light energy input*. Bioprocess and algae reactor technology, apoptosis. Berlin: Springer; 1998:123-52. doi: 10.1007/BFb0102298
 20. Carvalho AP, Meireles LA, Malcata FX. Microalgal reactors: a review of enclosed system designs and performances. *Biotechnol Prog* 2006;22(6):1490-506. doi: 10.1021/bp060065r
 21. Morweiser M, Kruse O, Hankamer B, Posten C. Developments and perspectives of photobioreactors for biofuel production. *Appl Microbiol Biotechnol* 2010;87(4):1291-301. doi: 10.1007/s00253-010-2697-x
 22. Chisti Y. Biodiesel from microalgae beats bioethanol. *Trends Biotechnol* 2008;26(3):126-31. doi: 10.1016/j.tibtech.2007.12.002
 23. Lam MK, Lee KT. Potential of using organic fertilizer to cultivate *Chlorella vulgaris* for biodiesel production. *Appl Energy* 2012;94:303-8. doi: 10.1016/j.apenergy.2012.01.075
 24. Dauta A, Devaux J, Piquemal F, Boumnick L. Growth rate of four freshwater algae in relation to light and temperature. *Hydrobiologia* 1990;207(1):221-6. doi: 10.1007/bf00041459
 25. Haggkvist K, Andersson C, Taesler R. Swedish meteorological and hydrological institute. Numerical Simulation of Fluid Flow and Heat/Mass Transfer Processes 2012;18:122.
 26. Jacob-Lopes E, Scoparo CHG, Lacerda LMCF, Franco TT. Effect of light cycles (night/day) on CO₂ fixation and biomass production by microalgae in photobioreactors. *Chem Eng Process* 2009;48(1):306-10. doi: 10.1016/j.cep.2008.04.007
 27. Amini Khoeyi Z, Seyfabadi J, Ramezanpour Z. Effect of light intensity and photoperiod on biomass and fatty acid composition of the microalgae, *Chlorella vulgaris*. *Aquac Int* 2012;20(1):41-9. doi: 10.1007/s10499-011-9440-1
 28. Choi YK, Kim HJ, Kumaran RS, Song HJ, Song KG, Kim KJ, et al. Enhanced growth and total fatty acid production of microalgae under various lighting conditions induced by flashing light. *Eng Life Sci* 2017;17(9):976-80. doi: 10.1002/elsc.201700001
 29. Zhao Y, Wang J, Zhang H, Yan C, Zhang Y. Effects of various LED light wavelengths and intensities on microalgae-based simultaneous biogas upgrading and digestate nutrient reduction process. *Bioresour Technol* 2013;136:461-8. doi: 10.1016/j.biortech.2013.03.051
 30. Xie T, Xia Y, Zeng Y, Li X, Zhang Y. Nitrate concentration-shift cultivation to enhance protein content of heterotrophic microalga *Chlorella vulgaris*: Over-compensation strategy. *Bioresour Technol* 2017;233:247-55. doi: 10.1016/j.biortech.2017.02.099
 31. Pietryczuk A, Czerpak R. Effect of traumatic acid on antioxidant activity in *Chlorella vulgaris* (Chlorophyceae). *Plant Growth Regul* 2011;65(2):279-86. doi: 10.1007/s10725-011-9599-5
 32. Qian H, Xu X, Chen W, Jiang H, Jin Y, Liu W, et al. Allelochemical stress causes oxidative damage and inhibition of photosynthesis in *Chlorella vulgaris*. *Chemosphere* 2009;75(3):368-75. doi: 10.1016/j.chemosphere.2008.12.040
 33. Gao QT, Wong YS, Tam NFY. Antioxidant responses of different microalgal species to nonylphenol-induced oxidative stress. *J Appl Phycol* 2017;29(3):1317-29. doi: 10.1007/s10811-017-1065-y
 34. Grudziński W, Krzeminska I, Luchowski R, Nosalewicz A, Gruszecki WI. Strong-light-induced yellowing of green microalgae *Chlorella*: A study on molecular mechanisms of the acclimation response. *Algal Res* 2016;16:245-54. doi: 10.1016/j.algal.2016.03.021
 35. Estevez MS, Malanga G, Puntarulo S. Iron-dependent oxidative stress in *Chlorella vulgaris*. *Plant Sci* 2001;161(1):9-17. doi: 10.1016/S0168-9452(01)00364-8
 36. Li Q, Liang Z, Ge F, Xu Y, Yang L, Zeng H. Alleviating CTAC and Flu combined pollution damage in *Chlorella vulgaris* by exogenous nitric oxide. *Chemosphere* 2014;96:39-45. doi: 10.1016/j.chemosphere.2013.07.012
 37. Shen YF, Liu L, Gong YX, Zhu B, Liu GL, Wang GX. Potential toxic effect of trifloxystrobin on cellular microstructure, mRNA expression and antioxidant enzymes in *Chlorella vulgaris*. *Environ Toxicol Pharmacol* 2014;37(3):1040-7. doi: 10.1016/j.etap.2014.04.006
 38. Dewys WD, Begg C, Lavin PT, Band PR, Bennett JM, Bertino JR, et al. Prognostic effect of weight loss prior to chemotherapy in cancer patients. Eastern Cooperative Oncology Group. *Am J Med* 1980;69(4):491-7. doi: 10.1016/S0149-2918(05)80001-3
 39. Liu L, Zhu B, Wang GX. Azoxystrobin-induced excessive reactive oxygen species (ROS) production and inhibition of photosynthesis in the unicellular green algae *Chlorella vulgaris*. *Environ Sci Pollut Res Int* 2015;22(10):7766-75. doi: 10.1007/s11356-015-4121-7
 40. Morschett H, Freier L, Rohde J, Wiechert W, von Lieres E, Oldiges M. A framework for accelerated phototrophic bioprocess development: integration of parallelized microscale cultivation, laboratory automation and Kriging-assisted experimental design. *Biotechnol Biofuels* 2017;10:26. doi: 10.1186/s13068-017-0711-6
 41. Alketife AM, Judd S, Znad H. Synergistic effects and optimization of nitrogen and phosphorus concentrations on the growth and nutrient uptake of a freshwater *Chlorella vulgaris*. *Environ Technol* 2017;38(1):94-102. doi: 10.1080/09593330.2016.1186227
 42. Schreiber C, Behrendt D, Huber G, Pfaff C, Widzowski J, Ackermann B, et al. Growth of algal biomass in laboratory and in large-scale algal photobioreactors in the temperate climate of western Germany. *Bioresour Technol* 2017;234:140-9. doi: 10.1016/j.biortech.2017.03.028
 43. Ramirez-Lopez C, Chairez I, Fernandez-Linares L. A novel culture medium designed for the simultaneous enhancement of biomass and lipid production by *Chlorella vulgaris* UTEX 26. *Bioresour Technol* 2016;212:207-16. doi: 10.1016/j.biortech.2016.04.051
 44. Josephine A, Niveditha C, Radhika A, Shali AB, Kumar TS, Dharani G, et al. Analytical evaluation of different carbon sources and growth stimulators on the biomass and lipid production of *Chlorella vulgaris* – Implications for biofuels. *Biomass Bioenergy* 2015;75:170-9. doi: 10.1016/j.biombioe.2015.02.016
 45. Liu J, Song Y, Liu Y, Ruan R. Optimization of growth conditions toward two-stage cultivation for lipid production of *Chlorella vulgaris*. *Environ Prog Sustain*

- Energy 2015;34(6):1801-7. doi: 10.1002/ep.12162
46. Chu FF, Shen XF, Lam PKS, Zeng RJ. Optimization of CO₂ concentration and light intensity for biodiesel production by *Chlorella vulgaris* FACHB-1072 under nitrogen deficiency with phosphorus luxury uptake. *J Appl Phycol* 2014;26(4):1631-8. doi: 10.1007/s10811-013-0200-7
 47. Kong WB, Hua SF, Cao H, Mu YW, Yang H, Song H, et al. Optimization of mixotrophic medium components for biomass production and biochemical composition biosynthesis by *Chlorella vulgaris* using response surface methodology. *J Taiwan Inst Chem Eng* 2012;43(3):360-7. doi: 10.1016/j.jtice.2011.11.007
 48. Dubey KK, Kumar S, Dixit D, Kumar P, Kumar D, Jawed A, et al. Implication of Industrial Waste for Biomass and Lipid Production in *Chlorella minutissima* Under Autotrophic, Heterotrophic, and Mixotrophic Grown Conditions. *Appl Biochem Biotechnol* 2015;176(6):1581-95. doi: 10.1007/s12010-015-1663-6
 49. Cho DH, Ramanan R, Heo J, Shin DS, Oh HM, Kim HS. Influence of limiting factors on biomass and lipid productivities of axenic *Chlorella vulgaris* in photobioreactor under chemostat cultivation. *Bioresour Technol* 2016;211:367-73. doi: 10.1016/j.biortech.2016.03.109
 50. Gonzalez-Garcinuno A, Sanchez-Alvarez JM, Galan MA, Martin Del Valle EM. Understanding and optimizing the addition of phytohormones in the culture of microalgae for lipid production. *Biotechnol Prog* 2016;32(5):1203-11. doi: 10.1002/btpr.2312
 51. Sharma AK, Sahoo PK, Singhal S. Screening and optimization of culture media for *Chlorella* sp. as a raw material for biodiesel production. *Int J Pharma Bio Sci* 2015;6(3):B251-62.
 52. Ramanna L, Guldhe A, Rawat I, Bux F. The optimization of biomass and lipid yields of *Chlorella sorokiniana* when using wastewater supplemented with different nitrogen sources. *Bioresour Technol* 2014;168:127-35. doi: 10.1016/j.biortech.2014.03.064
 53. Lu BL, Li MX, Qi L. The effect of light on the growth and product accumulation of *Chlorella*. *Adv Mat Res* 2013;724-725:323-9.
 54. Cui H, Meng F, Li F, Wang Y. Application of sodium erythorbate to promote the growth of *Chlorella vulgaris*. *J Appl Phycol* 2017;29(3):1135-44. doi: 10.1007/s10811-016-1021-2
 55. de Jesus SS, Maciel Filho R. Potential of algal biofuel production in a hybrid photobioreactor. *Chem Eng Sci* 2017;171:282-92. doi: 10.1016/j.ces.2017.05.041
 56. Gupta PL, Choi HJ, Pawar RR, Jung SP, Lee SM. Enhanced biomass production through optimization of carbon source and utilization of wastewater as a nutrient source. *J Environ Manage* 2016;184(Pt 3):585-95. doi: 10.1016/j.jenvman.2016.10.018
 57. Morowvat MH, Ghasemi Y. Medium optimization by artificial neural networks for maximizing the triglyceride-rich lipids from biomass of *Chlorella vulgaris*. *Int J Pharm Clin Res* 2016;8(10):1414-7.
 58. Chen T, Liu J, Guo B, Ma X, Sun P, Liu B, et al. Light attenuates lipid accumulation while enhancing cell proliferation and starch synthesis in the glucose-fed oleaginous microalga *Chlorella zofingiensis*. *Sci Rep* 2015;5:1-10. doi: 10.1038/srep14936
 59. Yeh KL, Chang JS, Chen WM. Effect of light supply and carbon source on cell growth and cellular composition of a newly isolated microalga *Chlorella vulgaris* ESP-31. *Eng Life Sci* 2010;10(3):201-8. doi: 10.1002/elsc.200900116
 60. Mayo AW. Effects of temperature and pH on the kinetic growth of unialga *Chlorella vulgaris* cultures containing bacteria. *Water Environ Res* 1997;69(1):64-72. doi: 10.2175/106143097X125191
 61. Castellanos C. Batch and continuous studies of *Chlorella vulgaris* in photobioreactors [dissertation] Ontario, Canada: University of Western Ontario; 2013.
 62. Lustigman B, Lee LH, Khalil A. Effects of nickel and pH on the growth of *Chlorella vulgaris*. *Bull Environ Contam Toxicol* 1995;55(1):73-80. doi: 10.1007/bf00212391
 63. Schenk PM, Thomas-Hall SR, Stephens E, Marx UC, Mussgnug JH, Posten C, et al. Second generation biofuels: high-efficiency microalgae for biodiesel production. *Bioenergy Res* 2008;1(1):20-43. doi: 10.1007/s12155-008-9008-8
 64. Widjaja A, Chien CC, Ju YH. Study of increasing lipid production from fresh water microalgae *Chlorella vulgaris*. *J Taiwan Inst Chem Eng* 2009;40(1):13-20. doi: 10.1016/j.jtice.2008.07.007
 65. Chen W, Zhang C, Song L, Sommerfeld M, Hu Q. A high throughput Nile red method for quantitative measurement of neutral lipids in microalgae. *J Microbiol Methods* 2009;77(1):41-7. doi: 10.1016/j.mimet.2009.01.001
 66. Lee JY, Yoo C, Jun SY, Ahn CY, Oh HM. Comparison of several methods for effective lipid extraction from microalgae. *Bioresour Technol* 2010;101 Suppl 1:S75-7. doi: 10.1016/j.biortech.2009.03.058
 67. Yoo C, Jun SY, Lee JY, Ahn CY, Oh HM. Selection of microalgae for lipid production under high levels carbon dioxide. *Bioresour Technol* 2010;101 Suppl 1:S71-4. doi: 10.1016/j.biortech.2009.03.030
 68. Starr RC. The culture collection of algae at the university of texas at austin. *J Phycol* 1978;14(Suppl 4):47-100. doi: 10.1111/j.1529-8817.1978.tb02507.x
 69. Keffer JE, Kleinheinz GT. Use of *Chlorella vulgaris* for CO₂ mitigation in a photobioreactor. *J Ind Microbiol Biotechnol* 2002;29(5):275-80. doi: 10.1038/sj.jim.7000313
 70. Alyabyev AJ, Loseva NL, Gordon LK, Andreyeva IN, Rachimova GG, Tribunskih VI, et al. The effect of changes in salinity on the energy yielding processes of *Chlorella vulgaris* and *Dunaliella maritima* cells. *Thermochim Acta* 2007;458(1):65-70. doi: 10.1016/j.tca.2007.03.003
 71. Abdel-Rahman MHM, Ali RM, Said HA. Alleviation of NaCl-induced effects on *Chlorella vulgaris* and *Chlorococcum humicola* by riboflavin application. *Int J Agric Biol* 2005;7:58-62.
 72. Richmond A, Karg S, Boussiba S. Effects of Bicarbonate and Carbon Dioxide on the Competition between *Chlorella vulgaris* and *Spirulina platensis*. *Plant Cell Physiol* 1982;23(8):1411-7. doi: 10.1093/oxfordjournals.pcp.a076489
 73. Zheng H, Gao Z, Zhang Q, Huang H, Ji X, Sun H, et al. [Effect of inorganic carbon source on lipid production with autotrophic *Chlorella vulgaris*]. *Sheng Wu Gong Cheng Xue Bao* 2011;27(3):436-44.
 74. Sirisansaneeyakul S, Singhasuwan S, Choorit W, Phoopat N, Garcia JL, Chisti Y. Photoautotrophic production of

- lipids by some *Chlorella* strains. *Mar Biotechnol* (NY) 2011;13(5):928-41. doi: 10.1007/s10126-010-9355-2
75. Kumar K, Das D. Growth characteristics of *Chlorella sorokiniana* in airlift and bubble column photobioreactors. *Bioresour Technol* 2012;116:307-13. doi: 10.1016/j.biortech.2012.03.074
 76. Aishvarya V, Pradhan N, Nayak RR, Sukla LB, Mishra BK. Enhanced inorganic carbon uptake by *Chlorella* sp. IMMTCC-2 under autotrophic conditions for lipid production and CO₂ sequestration. *J Appl Phycol* 2012;24(6):1455-63. doi: 10.1007/s10811-012-9801-9
 77. Devgawami CR, Kalita MC, Talukdar J, Bora R, Sharma P. Studies on the growth behavior of *Chlorella*, *Haematococcus* and *Scenedesmus* sp. in culture media with different concentrations of sodium bicarbonate and carbon dioxide gas. *Afr J Biotechnol* 2011;10(61):13128-38.
 78. Sun X, Zhong Y, Huang Z, Yang Y. Selenium accumulation in unicellular green alga *Chlorella vulgaris* and its effects on antioxidant enzymes and content of photosynthetic pigments. *PLoS One* 2014;9(11):e112270. doi: 10.1371/journal.pone.0112270
 79. Goiris K, Muylaert K, De Cooman L. Microalgae as a novel source of antioxidants for nutritional applications. In: Kim SK, ed. *Handbook of marine microalgae*. Boston: Academic Press; 2015:269-80. doi: 10.1016/B978-0-12-800776-1.00017-0
 80. Barsanti L, Coltelli P, Evangelista V, Frassanito AM, Passarelli V, Vesentini N, et al. Oddities and curiosities in the algal world. In: Evangelista V, Barsanti L, Frassanito AM, eds. *Algal toxins: nature, occurrence, effect and detection*. Springer; 2008:353-91.
 81. Skjanes K, Rebours C, Lindblad P. Potential for green microalgae to produce hydrogen, pharmaceuticals and other high value products in a combined process. *Crit Rev Biotechnol* 2013;33(2):172-215. doi: 10.3109/07388551.2012.681625
 82. Takahashi S, Badger MR. Photoprotection in plants: a new light on photosystem II damage. *Trends Plant Sci* 2011;16(1):53-60. doi: 10.1016/j.tplants.2010.10.001
 83. Malanga G, Calmanovici G, Puntarulo S. Oxidative damage to chloroplasts from *Chlorella vulgaris* exposed to ultraviolet-B radiation. *Physiol Plant* 1997;101(3):455-62. doi: 10.1111/j.1399-3054.1997.tb01023.x
 84. Devos N, Ingouff M, Loppes R, Matagne RF. RUBISCO adaptation to low temperatures: a comparative study in psychrophilic and mesophilic unicellular algae. *J Phycol* 1998;34(4):655-60. doi: 10.1046/j.1529-8817.1998.340655.x
 85. Tjahjono AE, Hayama Y, Kakizono T, Terada Y, Nishio N, Nagai S. Hyper-accumulation of astaxanthin in a green alga *Haematococcus pluvialis* at elevated temperatures. *Biotechnol Lett* 1994;16(2):133-8. doi: 10.1007/bf01021659
 86. Miranda MS, Sato S, Mancini-Filho J. Antioxidant activity of the microalga *Chlorella vulgaris* cultured on special conditions. *Boll Chim Farm* 2001;140(3):165-8.
 87. Stoyanova-Bakalova E, Toncheva-Panova T. Subcellular adaptation to salinity and irradiance in *Dunaliella salina*. *Biol Plant* 2003;47(2):233-6. doi: 10.1023/B:BIOP.0000022257.34916.c5
 88. Oren A. Diversity of organic osmotic compounds and osmotic adaptation in cyanobacteria and algae. In: Seckbach J, ed. *Algae and cyanobacteria in extreme environments*. Springer; 2007:639-55. doi: 10.1007/978-1-4020-6112-7_35
 89. Hadi MR, Shariati M, Afsharzadeh S. Microalgal biotechnology: Carotenoid and glycerol production by the green algae *Dunaliella* isolated from the Gave-Khooni salt marsh, Iran. *Biotechnol Bioprocess Eng* 2008;13(5):540. doi: 10.1007/s12257-007-0185-7
 90. Rao AR, Dayananda C, Sarada R, Shamala TR, Ravishankar GA. Effect of salinity on growth of green alga *Botryococcus braunii* and its constituents. *Bioresour Technol* 2007;98(3):560-4. doi: 10.1016/j.biortech.2006.02.007
 91. Pandit PR, Fulekar MH, Karuna MSL. Effect of salinity stress on growth, lipid productivity, fatty acid composition, and biodiesel properties in *Acutodesmus obliquus* and *Chlorella vulgaris*. *Environ Sci Pollut Res Int* 2017;24(15):13437-51. doi: 10.1007/s11356-017-8875-y
 92. Coesel SN, Baumgartner AC, Teles LM, Ramos AA, Henriques NM, Cancela L, et al. Nutrient limitation is the main regulatory factor for carotenoid accumulation and for PsII and PsI steady state transcript levels in *Dunaliella salina* (Chlorophyta) exposed to high light and salt stress. *Mar Biotechnol* (NY) 2008;10(5):602-11. doi: 10.1007/s10126-008-9100-2
 93. He P, Duncan J, Barber J. Astaxanthin Accumulation in the Green Alga *Haematococcus pluvialis*: Effects of Cultivation Parameters. *J Integr Plant Biol* 2007;49(4):447-51. doi: 10.1111/j.1744-7909.2007.00468.x
 94. Griffiths MJ, Harrison STL. Lipid productivity as a key characteristic for choosing algal species for biodiesel production. *J Appl Phycol* 2009;21(5):493-507. doi: 10.1007/s10811-008-9392-7
 95. Solovchenko AE, Khozin-Goldberg I, Didi-Cohen S, Cohen Z, Merzlyak MN. Effects of light intensity and nitrogen starvation on growth, total fatty acids and arachidonic acid in the green microalga *Parietochloris incisa*. *J Appl Phycol* 2008;20(3):245-51. doi: 10.1007/s10811-007-9233-0
 96. Durmaz Y. Vitamin E (α -tocopherol) production by the marine microalgae *Nannochloropsis oculata* (Eustigmatophyceae) in nitrogen limitation. *Aquaculture* 2007;272(1):717-22. doi: 10.1016/j.aquaculture.2007.07.213
 97. Sakarika M, Kornaros M. Kinetics of growth and lipids accumulation in *Chlorella vulgaris* during batch heterotrophic cultivation: Effect of different nutrient limitation strategies. *Bioresour Technol* 2017;243:356-65. doi: 10.1016/j.biortech.2017.06.110
 98. Brennan L, Owende P. Biofuels from microalgae—a review of technologies for production, processing, and extractions of biofuels and co-products. *Renew Sust Energy Rev* 2010;14(2):557-77. doi: 10.1016/j.rser.2009.10.009
 99. Sheehan J, Dunahay T, Benemann J, Roessler P. *Look back at the US department of energy's aquatic species program: biodiesel from algae*. National Renewable Energy Laboratory; 1998:328. doi: 10.2172/15003040
 100. Gerardi MH. *The microbiology of anaerobic digesters*. John Wiley & Sons; 2003. doi: 10.1002/0471468967